

## Diabetic ketoacidosis

### Definition

Diabetic ketoacidosis (DKA) is a life-threatening problem that affects people with diabetes. It occurs when the body starts breaking down fat at a rate that is much too fast. The liver processes the fat into a fuel called ketones, which causes the blood to become acidic.

### Causes


DKA happens when the signal from insulin in the body is so low that:

1. Glucose (blood sugar) can't go into cells to be used as a fuel source.
2. The liver makes a huge amount of blood sugar.
3. Fat is broken down too rapidly for the body to process.

The fat is broken down by the liver into a fuel called ketones. Ketones are normally produced by the liver when the body breaks down fat after it has been a long time since your last meal. These ketones are normally used by the muscles and the heart. When ketones are produced too quickly and build up in the blood, they can be toxic by making the blood acidic. This condition is known as ketoacidosis.

DKA is sometimes the first sign of type 1 diabetes in people who have not yet been diagnosed. It can also occur in someone who has already been diagnosed with type 1 diabetes. Infection, injury, a serious illness, missing doses of insulin shots, or the stress of surgery can lead to DKA in people with type 1 diabetes.

People with type 2 diabetes can also develop DKA, but it is less common and less severe. It is usually triggered by prolonged uncontrolled blood sugar



ADAM.

### Oral Glucose Tolerance Test



No food or drink 8 to 12 hours prior to test



Drink glucose



Blood is tested two hours later

High glucose level = potential diabetes

ADAM.

severe illness or infection.

## Symptoms

Common symptoms of DKA can include:

- Decreased alertness
- Deep, rapid breathing
- Dehydration
- Dry skin and mouth
- Flushed face
- Frequent urination or thirst that lasts for a day or more
- Fruity-smelling breath
- Headache
- Muscle stiffness or aches
- Nausea and vomiting
- Stomach pain



An insulin pump can be worn discretely under clothing as it administers insulin to the person with diabetes



ADAM.

## Exams and Tests

Ketone testing may be used in type 1 diabetes to screen for early ketoacidosis. The ketone test is usually done using a urine sample or a blood sample.

Ketone testing is usually done when DKA is suspected:

- Most often, urine testing is done first.
- If the urine is positive for ketones, most often a ketone called beta-hydroxybutyrate is measured in the blood. This is the most common ketone measured. The other main ketone is acetoacetate.

Other tests for ketoacidosis include:

- Arterial blood gas
- Basic metabolic panel, (a group of blood tests that measure your sodium and potassium levels, kidney function, and other chemicals and functions, including the anion gap)
- Blood glucose test
- Blood pressure measurement
- Osmolality blood test

## Treatment

The goal of treatment is to correct the high blood sugar level with insulin. Another goal is to replace fluids lost through urination, loss of appetite, and vomiting if you have these symptoms.

If you have diabetes, it is likely your health care provider told you how to spot the warning signs of DKA. If you think you have DKA, test for ketones using urine strips. Some glucose meters can also measure blood ketones. If ketones are present, call your provider right away. DO NOT delay. Follow any instructions you are given.

treatment for DKA. Then providers will also search for and treat the cause of DKA, such as an infection.

## Outlook (Prognosis)

Most people respond to treatment within 24 hours. Sometimes, it takes longer to recover.

If DKA is not treated, it can lead to severe illness or death.

## Possible Complications

Health problems that may result from DKA include any of the following:

- Fluid buildup in the brain (cerebral edema)
- Heart stops working (cardiac arrest)
- Kidney failure

## When to Contact a Medical Professional

DKA is often a medical emergency. Call your provider if you notice symptoms of DKA.

Go to the emergency room or call the local emergency number (such as 911) if you or a family member with diabetes has any of the following:

- Decreased consciousness
- Fruity breath
- Nausea and vomiting
- Trouble breathing

## Prevention

If you have diabetes, learn to recognize the signs and symptoms of DKA. Know when to test for ketones, such as when you are sick.

If you use an insulin pump, check often to see that insulin is flowing through the tubing. Make sure the tube is not blocked, kinked or disconnected from the pump.

## Alternative Names

DKA; Ketoacidosis; Diabetes - ketoacidosis

---

Review Date: January 26, 2020.

Reviewed By: Brent Wisse, MD, board certified in Metabolism/Endocrinology, Seattle, WA. Internal review and update on 06/04/2021 by David Zieve, MD, MHA, Medical Director, Brenda Conaway, Editorial Director, and the A.D.A.M. Editorial team. Editorial update 09/30/2021.

A.D.A.M., Inc. is accredited by URAC, for Health Content Provider ([www.uran.org](http://www.uran.org)). URAC's [accreditation program](#) is an independent audit to verify that A.D.A.M. follows rigorous standards of quality and accountability. A.D.A.M. is among the first to achieve this important distinction for online health information and services. Learn more about A.D.A.M.'s [editorial policy](#), [editorial process](#) and [privacy policy](#). A.D.A.M. is also a founding member of Hi-Ethics. This site complies with the HONcode standard for trustworthy health information: [verify here](#).

The information provided herein should not be used during any medical emergency or for the diagnosis or treatment of any medical condition. A licensed medical professional should be consulted for diagnosis and treatment of any and all medical conditions. Call 911 for all medical emergencies. Links to other sites are provided for information only -- they do not constitute endorsements of those other sites. © 1997-2022 A.D.A.M., Inc. Any duplication or distribution of the information contained herein is strictly prohibited.

